

EXAMPLES OF NON INTEGER DIMENSIONAL GEOMETRIES

R. TRINCHERO

ABSTRACT. Two examples of spectral triples with non-integer dimension spectrum are considered. These triples involve commutative C^* -algebras. The first example has complex dimension spectrum and trivial differential algebra. The other is a parameter dependent deformation of the canonical spectral triple over S^1 . Its dimension spectrum includes real non-integer values. It has a non-trivial differential algebra and in contrast with the one dimensional case there are no junk forms for a non-vanishing deformation parameter. The distance on this space depends non-trivially on this parameter.

1. INTRODUCTION

The dimension of a space is a basic concept of particular relevance both in nature and in mathematics. Non-commutative geometry¹ provides a generalization of classical geometry. It is the object of this work to study some examples of non-classical geometrical objects within this framework. In particular, examples where the non-commutative analog of dimension for the corresponding spaces can take non-integer values are dealt with. The study of these geometries contributes both to the implementation of physical ideas in these spaces and to the exploration in concrete examples of the basic concepts in non-commutative geometry.

In the realm of quantum field theory, the widely employed dimensional regularization technique[2] provides a hint that such non-integer dimensional spaces could be of relevance in physics. This study intends to go a step further in realizing those spaces as concrete geometrical objects. In the field theoretic setting a desirable outcome in the direction of the ideas mentioned above would be, for example, to obtain an implementation of dimensional regularization at a Lagrangian level, or, in a more general setting, to really consider the possibility that physical space is non-integer dimensional.

The general picture of non-commutative geometry shows that there are many ways to construct spaces with non integer dimensions. In this work we look for, so to say, the simplest and easiest to treat examples of such spaces. A first step in this direction is to realize that it is possible to get non-integer dimensional geometries even when the underlying C^* -algebra of the spectral triple is commutative. The examples presented below are of this kind.

¹The reader is referred to [1] for an account of this theory.

The generalization of the concept of dimension in non-commutative geometry is provided by the Connes-Moscovici definition of dimension spectrum[3]. Section 2 presents an example with complex dimension spectrum but trivial differential algebra. It can be considered as a simple example focused on getting a non-trivial dimension spectrum. Section 3 presents a deformation of the canonical spectral triple² whose dimension spectrum includes real non-integer values. The main results for this last example obtained in this work are related to junk forms and distance. As mentioned in the abstract, no junk forms appear for non-vanishing values of the deformation parameter α . This fact leads to a differential algebra which involves forms of arbitrary degree and is neither anti-commutative nor commutative. This situation may be compared with the canonical case where junk forms play a crucial role in getting the usual differential algebra. This important qualitative difference between the integer dimensional case and the deformed one is, in a certain vague sense, analog to the absence of poles in dimensionally regularized Feynman integrals.

Regarding distance, this turns out to be a function of the coordinates and α . This function involves a $\Gamma(\varepsilon)(\varepsilon = 1 - \alpha)$ which does not come, as in dimensional regularization, from an analytical continuation of the factorial appearing in the solid angle integral. In fact, there is no solid angle involved since we start from a support space for the commutative C^* -algebra in the spectral triple which is one dimensional.

2. EXAMPLE WITH COMPLEX DIMENSION SPECTRUM.

2.1. General setting. Here we deal with the triple $(\mathcal{A}, \mathcal{H}, D)$ where,

2.1.1. Algebra. $\mathcal{A}$ is the $\star$ -algebra generated by $1, a, \bar{a}$ with the following multiplication rules,

$$(1) \quad 1 \cdot a = a \cdot 1 = 1, 1 \cdot \bar{a} = \bar{a} \cdot 1 = 1, a \cdot \bar{a} = \bar{a} \cdot a$$

where the star structure is given by,

$$(2) \quad a^\star = \bar{a}, 1^\star = 1$$

This algebra is isomorphic to the one of polynomials in a complex variable z and its complex conjugate $\bar{z}$. The norm for this algebra will be given below in terms of a representation in a Hilbert space $\mathcal{H}$.

2.1.2. Hilbert space. $\mathcal{H} = L^2(S_{AP}^1)$ = square integrable anti-periodic functions on S^1 , thus³,

$$(3) \quad \psi(x), 0 \leq x \leq 2\pi \in \mathcal{H} \Rightarrow \psi(x) = -\psi(x + 2\pi)$$

²The contents of section 2 and subsections 3.1 and 3.2 appeared first in reference [4].

³This space can be identified with the one of square integrable sections of the associated spinor bundle corresponding to the anti-periodic spin structure over S^1 .

with the scalar product given by,

$$(4) \quad \langle \psi | \varphi \rangle = \frac{1}{2\pi} \int_0^{2\pi} \psi^*(x) \varphi(x)$$

a basis for this Hilbert space is given by the functions,

$$(5) \quad \langle x | k \rangle = e^{i(k+1/2)x}, \quad k \in \mathbb{Z}$$

this basis is orthonormal,

$$(6) \quad \begin{aligned} \langle k | m \rangle &= \int_x \langle k | x \rangle \langle x | m \rangle = \frac{1}{2\pi} \int_0^{2\pi} dx e^{-i(k+1/2)x} e^{i(m+1/2)x} \\ &= \delta_{km} \end{aligned}$$

2.1.3. *Representation of $\mathcal{A}$ in $\mathcal{H}$.* The following representation $\pi : \mathcal{A} \rightarrow \mathcal{H}$ is considered,

$$(7) \quad \pi(a) = \sum_k |k + 1/2|^\varepsilon |k\rangle \langle k|, \quad \varepsilon \in \mathbb{C}, |\varepsilon| < 1, \Re(\varepsilon) > 0$$

this definition has the property,

$$(8) \quad \pi(a^n) = \pi(a)^n$$

the representant of the star element is defined by,

$$(9) \quad \pi(\bar{a}) = \pi(a)^\dagger = \sum_k |k + 1/2|^{\bar{\varepsilon}} |k\rangle \langle k|$$

where $\bar{\varepsilon}$ is the complex conjugate of ε . The norm of the eigenvalues of $\pi(a)$ are always bounded as can be seen from the following equation,

$$(10) \quad ||k + 1/2|^{-\varepsilon}| = e^{-\Re(\varepsilon) \ln |k+1/2|}$$

and the requirement that $\Re(\varepsilon) > 0$. Thus the representative of any element in the algebra is bounded. Defining the norm for the algebra in terms of the operator norm in this representation leads to a pre- C^* -algebra structure in $\mathcal{A}$. Next this pre- C^* -algebra structure is completed to a C^* -algebra structure in $\mathcal{A}$.

2.1.4. *The Dirac operator.* The Dirac operator is taken to be,

$$(11) \quad D = -i\partial$$

this operator is selfadjoint in $\mathcal{H}$, and its resolvent is compact. Furthermore,

$$(12) \quad [D, \pi(a)] = 0, \quad \forall a \in \mathcal{A}$$

which is bounded. Thus the triple fulfills all the properties required for it to be a spectral triple.

2.2. Dimension spectrum of the triple. The definition of dimension spectrum of a spectral triple is briefly reviewed. Let δ denote the derivation $\delta : L(\mathcal{H}) \rightarrow L(\mathcal{H})$ defined by,

$$(13) \quad \delta(T) = [|D|, T] \quad , T \in L(\mathcal{H})$$

Let $\mathcal{B}$ denote the algebra generated by the elements,

$$(14) \quad \delta^n(\pi(a)), \ a \in \mathcal{A}, \ n \geq 0 \ (\delta^0(\pi(a)) = \pi(a))$$

Definition 2.1. [Connes-Moscovici] *Discrete dimension spectrum.* A spectral triple has discrete dimension spectrum Sd if $Sd \subset \mathbb{C}$ is discrete and for any element $b \in \mathcal{B}$ the function,

$$(15) \quad \zeta_b(z) = Tr[\pi(b) |D|^{-z}]$$

extends holomorphically to $\mathbb{C}/Sd$.

This definition is given in the paper [3], where the additional assumption is made that,

$$(16) \quad \mathcal{A} \subset \cap_{n>0} Dom \delta^n \text{ and } [D, \mathcal{A}] \subset \cap_{n>0} Dom \delta^n$$

which clearly holds for this case since $\mathcal{B} = Imag(\delta^0) = \mathcal{A}$. Thus the following generic ζ -function is considered,

$$(17) \quad \zeta_{P(a, \bar{a})}(z) = Tr[\pi(P(a, \bar{a})) | -i\partial|^{-z}]$$

where $P(a, \bar{a})$ is a polynomial in a and $\bar{a}$ given by,

$$(18) \quad P(a, \bar{a}) = \sum_{n, m \in \mathbb{N}+0} p_{nm} a^n \bar{a}^m$$

then,

$$\begin{aligned} \zeta_{P(a, \bar{a})}(z) &= \sum_{k \in \mathbb{Z}} \langle \psi_k | P(|-i\partial|^{-\varepsilon}, |-i\partial|^{-\bar{\varepsilon}}) | \psi_k \rangle |k + 1/2|^{-z} \\ &= 2 \sum_{k \in \mathbb{N}+0} P(|k + 1/2|^{-\varepsilon}, |k + 1/2|^{-\bar{\varepsilon}}) |k + 1/2|^{-z} \\ &= 2 \sum_{n, m \in \mathbb{N}+0} p_{nm} 2^{z+n\varepsilon+m\bar{\varepsilon}} \sum_{k \in \mathbb{N}+0} |2k + 1|^{-n\varepsilon-m\bar{\varepsilon}-z} \\ (19) \quad &= 2 \sum_{n, m \in \mathbb{N}+0} p_{nm} (2^{z+n\varepsilon+m\bar{\varepsilon}} - 1) \zeta_R(z + n\varepsilon + m\bar{\varepsilon}) \end{aligned}$$

where $\zeta_R(z)$ denotes the Riemann ζ -function, which has a simple pole at $z = 1$. Therefore the poles of $\zeta_{P(a, \bar{a})}(z)$ are located at,

$$(20) \quad z = 1 - n\varepsilon - m\bar{\varepsilon} \ , n, m \in \mathbb{N} + 0$$

Eq. 20 gives the dimension spectrum for this space. This spectrum is plotted in Fig.1 for $\varepsilon = 1 + i/2$.




FIGURE 1. Discrete dimension spectrum for the triple of sec. 2 with $\varepsilon = 1 + i/2$.

3. DEFORMATION OF THE CANONICAL SPECTRAL TRIPLE ON S^1

3.1. General setting. Here we deal with the spectral triple $(\mathcal{A}, \mathcal{H}, D)$ where,

3.1.1. Algebra. $\mathcal{A} = F(S^1)$ is the commutative C^* -algebra of smooth functions over S^1 . This algebra can be characterized by generators, $b_n, n \in \mathbb{Z}$ satisfying the relations,

$$b_n b_m = b_{n+m}, b_n^* = b_{-n}$$

3.1.2. Hilbert space. $\mathcal{H} = L^2(S^1_{AP})$ is the Hilbert space of square integrable anti-periodic functions over S^1 , basis $|k\rangle, k \in \mathbb{Z}$, such that

$$(21) \quad \langle x | k \rangle = e^{i(k+1/2)x}, n \in \mathbb{Z}, x \in [0, 2\pi] \text{ and } \langle l | k \rangle = \delta_{lk}$$

3.1.3. Representation of $\mathcal{A}$ in $\mathcal{H}$.

$$(22) \quad \pi(b_n) = \sum_{k \in \mathbb{Z}} |k+n\rangle \langle k|$$

3.1.4. The Dirac operator. $D \in L(\mathcal{H})$ is given by,

$$(23) \quad D = \sum_{k \in \mathbb{Z}} f_\alpha(k) |k\rangle \langle k|$$

The following conditions on the functions $f_\alpha(k)$ are required ,

- $\lim_{\alpha \rightarrow 0} f_\alpha(k) = k$. This condition ensures that in this limit the canonical spectral triple is recovered.
- $\exists C > 0 : |f_\alpha(k+n) - f_\alpha(k)| < C \ \forall k, n$. This implies that $[D, \pi(k)]$ is bounded in $L(\mathcal{H})$.
- $f_\alpha(k) = f_\alpha(k)^*$ implies that D is selfadjoint.

- $\lim_{k \rightarrow \infty} \frac{1}{f_\alpha(k) - \lambda} \rightarrow 0, \quad \forall \lambda \in \mathbb{R}$ implies that D has Compact resolvent.

The following choice of $f_\alpha(k)$ will be considered from now on,

$$(24) \quad f_\alpha(k) = \frac{(k + 1/2)^\varepsilon + [(k + 1/2)^\varepsilon]^*}{2}$$

where $\varepsilon \in \mathbb{R}$, $\varepsilon = 1 - \alpha$, $\alpha > 0$ and the power is defined by the principal branch of the logarithm, i.e.,

$$(25) \quad (k + 1/2)^\varepsilon = e^{\varepsilon \ln |k+1/2| + i\varepsilon \arg(k+1/2)}$$

where $\arg(k + 1/2) = 0$ or π . This implies that the power in (25) is real only for integer ε . This fact explains why the complex conjugate is summed up in (24). Using (25) it follows that,

$$(26) \quad f_\alpha(k) = \begin{cases} |k + 1/2|^\varepsilon, & k \geq 0 \\ |k + 1/2|^\varepsilon \cos(\pi\varepsilon), & k < 0 \end{cases}$$

This choice fulfills all the conditions required above.

3.2. Dimension spectrum of the triple. The property (16) holds for the choice (24). According to definition 2.1 the discrete dimension spectrum is the union of the poles in $\zeta_b(z)$ for any $b \in \mathcal{B}$. Each of such poles can be considered to describe the dimension of a certain piece of the whole space. If we take elements b in the image of δ^0 , i.e. $b \in \mathcal{A}$, the following zeta function is obtained,

$$(27) \quad \zeta_{b_n}(z) = \delta_{n,0} \sum_k |f_\alpha(k)|^{-z}$$

For the choice (24) the last equation leads to,

$$\begin{aligned} \zeta_{b_n}(z) &= \delta_{n,0} \left(\sum_{k=0}^{\infty} |k + 1/2|^{-\varepsilon z} + \sum_{k=-1}^{-\infty} |k + 1/2|^{-\varepsilon z} \cos(\pi\varepsilon) \right) \\ &= \delta_{n,0} [1 + \cos(\pi\varepsilon)] \sum_{k=0}^{\infty} |k + 1/2|^{-\varepsilon z} \\ (28) \quad &= \delta_{n,0} [1 + \cos(\pi\varepsilon)] (2^{\varepsilon z} - 1) \zeta_R(\varepsilon z) \end{aligned}$$

where $\zeta_R(z)$ denotes the Riemann zeta function. Thus $\zeta_{b_n}(z)$ has a simple isolated pole at,

$$(29) \quad z = 1/\varepsilon$$

Since we are simply searching for a deformation of the canonical triple with non-integer dimension we shall concentrate on this particular piece of this "space".

3.3. Differential algebra. Basic definition,

$$(30) \quad \pi(db_n) = [D, \pi(b_n)] = \sum_l [f_\alpha(l+n) - f_\alpha(l)] |l+n\rangle\langle l|$$

We look for junk forms, i.e., forms ω such that,

$$(31) \quad \pi(\omega) = 0 \quad , \pi(d\omega) \neq 0$$

- For the case of ω a zero form, the only solution of $\pi(\omega) = 0$ is $\omega = 0$, and $\pi(d\omega) = 0$ if $\omega = 0$. Thus in this case there are no junk 1-forms.
- For the case of ω a generic one form,

$$\begin{aligned} \pi(\omega) &= \pi\left(\sum_{m,n} \alpha_{mn} b_m db_n\right) \\ &= \sum_{m,n,l} \alpha_{mn} (f_\alpha(l+n) - f_\alpha(l)) |l+m+n\rangle\langle l| \\ (32) \quad &= \sum_{r,n,l} \alpha_{r-n,n} (f_\alpha(l+n) - f_\alpha(l)) |l+r\rangle\langle l| \end{aligned}$$

Now the operators $|r\rangle\langle s|$ and $|p\rangle\langle q|$ are linearly independent if $r \neq p$ and/or $s \neq q$. Thus there can only be cancellations in the r.h.s. of (32) between terms with $r = m + n$ and l fixed. Hence the vanishing of $\pi(\omega)$ implies,

$$(33) \quad \sum_n v_n^{(r)} w^{(l)}(n) = 0 \quad , \forall r, l$$

where,

$$(34) \quad v_n^{(r)} = \alpha_{r-n,n} \quad , w^{(l)}(n) = f_\alpha(l+n) - f_\alpha(l)$$

Next, the differential of such 1-form ω with $\pi(\omega) = 0$ is considered,

$$\begin{aligned} \pi(d\omega) &= \sum_{m,n} \alpha_{mn} \pi(db_m db_n) \\ &= \sum_{m,n,k,l} \alpha_{mn} w^{(k)}(m) w^{(l)}(n) |k+m\rangle\langle k| |l+n\rangle\langle l| \\ &= \sum_{m,n,l} \alpha_{mn} w^{(l+n)}(m) w^{(l)}(n) |l+m+n\rangle\langle l| \\ (35) \quad &= \sum_{r,n,l} \alpha_{r-n,n} w^{(l+n)}(r-n) w^{(l)}(n) |l+r\rangle\langle l| \end{aligned}$$

below the issue of junk forms is studied both for the canonical and the deformed spectral triples. The following review of the canonical case will hopefully make clearer the difference with the deformed case.

1. $\alpha = \mathbf{0}$ ($f_0(k) = k + 1/2$). Eqs. (33) and (34) leads to,

$$(36) \quad 0 = \pi(\omega) = \sum_{m,n,l} \alpha_{m,n} n |l+m+n\rangle\langle l|$$

So, the equation to solve is,

$$(37) \quad 0 = \sum_n v_n^{(r)} n = 0, \forall r$$

Two ways of dealing with this equation will be considered:

(i) First we consider the equation,

$$(38) \quad (\mathbf{v}, \mathbf{N}) = 0$$

where $\mathbf{N}$ is the vector $(-N, -N+1, \dots, -1, 0, 1, \dots, N)$ in a vector space $\mathbb{C}^{2N+1}$, $(,)$ is the canonical scalar product in that space and the equation should be solved for the vector $\mathbf{v}$, i.e. we look for all vectors orthogonal to $\mathbf{N}$. Therefore, there are $2N$ independent solutions to this equation, which can be chosen to be the following,

- (a) For each $I = 1, \dots, N$, $v_I = 1 = v_{-I}$ the rest zero are N solutions.
- (b) $v_0 = 1$ the rest zero is solution.
- (c) For $I = 2, \dots, N$, $v_1 = I = -v_{-1}$, $v_I = -1 = -v_{-I}$ the rest zero are $N-1$ solutions.

these are a set of $2N$ independent solutions, any other solution can be expressed as a linear combination of these ones. Thus, the solutions for α_{mn} are given for each $m+n=r$ fixed in terms of the ones above by,

$$(39) \quad \alpha_{r-n,n} = v_n^{(r)}$$

For different values of r , there can correspond different solutions for $v_n^{(r)}$.

(ii) Defining $v(x)$ by,

$$(40) \quad v^{(r)}(x) = \sum_n v_n^{(r)} e^{i(n+1/2)x}, 0 \leq x \leq 2\pi$$

eq.(37) can be written as,

$$(41) \quad -i \frac{d}{dx} \left(v^{(r)}(x) e^{-ix/2} \right) \Big|_{x=0} = 0 \quad \forall r$$

which implies that $v^{(r)}(x) e^{-ix/2}$ can be any periodic function with vanishing derivative at the origin. Note that the solutions appearing in 1., in terms of $v(x)$ are: (a) $v(x) = 2 e^{ix/2} \cos kx$, $k = 1, 2, \dots$, (b) $e^{ix/2}$ (c) $v(x) = 2i e^{ix/2} [k \sin(x) - \sin(kx)]$ $k = 2, 3, \dots$ whose linear combinations generate any function satisfying (41). Now, for the differential $\pi(d\omega)$ using (40) we obtain,

$$(42) \quad \pi(d\omega) = \sum_{r,n,l} \alpha_{r-n,n} (r-n) n |l+r\rangle \langle l|$$

which in terms of $v^{(r)}(x)$ can be written as,

$$(43) \quad \begin{aligned} \pi(d\omega) = & \sum_r \left\{ i \frac{d}{dx} \left[e^{-irx} - i \frac{d}{dx} (v^{(r)}(x) e^{-ix/2}) \right] \right\}_{x=0} \times \\ & \times \sum_l |l+r\rangle \langle l| \end{aligned}$$

the vanishing of the l.h.s. in (43) imply the vanishing of the expression between braces, which in turn implies, due to (41), that,

$$\left[\frac{d^2}{dx^2} (v^{(r)}(x) e^{-ix/2}) \right]_{x=0} = 0$$

thus since $\pi(\omega) = 0$ does not imply $\pi(d\omega) = 0$, there are junk forms in this case. Indeed one can show that any two form is a junk form in this case. ⁴

2. $\alpha \neq 0$. It is shown that,

Proposition 3.1. There are no non-trivial junk 1-forms for $\alpha \neq 0$.

Proof. Replacing (24) in (34) leads to,

$$(48) \quad \begin{aligned} w^{(l)}(n) &= f_\alpha(l+n) - f_\alpha(l) \\ &= \frac{(l+n+1/2)^\varepsilon + [(l+n+1/2)^\varepsilon]^* - (l+1/2)^\varepsilon - [(l+1/2)^\varepsilon]^*}{2} \end{aligned}$$

⁴To make contact with other approaches to this problem, the following 1-form is considered,

$$(44) \quad \omega_m = b_m db_m - db_m b_m$$

replacing (22) for $\pi(b_m)$ the following expression is obtained,

$$(45) \quad \pi(\omega_m) = \sum_l [f_\alpha(l+m) - f_\alpha(l) - f_\alpha(l+2m) + f_\alpha(l+m)] |l+2m\rangle \langle l|$$

which for the canonical case $f_\alpha(l) = l$ leads to $\pi(\omega_m) = 0$. However,

$$(46) \quad \begin{aligned} \pi(d\omega_m) &= \pi(db_m)\pi(db_m) \\ &= \sum_{k,l} [f_\alpha(k+m) - f_\alpha(k)] [f_\alpha(l+m) - f_\alpha(l)] |k+m\rangle \langle k| |l+m\rangle \langle l| \\ &= \sum_l [f_\alpha(l+2m) - f_\alpha(l+m)] [f_\alpha(l+m) - f_\alpha(l)] |l+2m\rangle \langle l| \\ &= \sum_l m^2 |l+2m\rangle \langle l| \end{aligned}$$

which does not vanish. Furthermore, one can show that any two form ω can be written as,

$$(47) \quad \omega = \sum_n f_n w_n$$

where f_n are elements of $\mathcal{A}$. Thus, leading to the result that in the canonical case any two form is a junk form.

Employing the binomial expansion for the first two terms in the last equation leads to,,

$$\begin{aligned}
 w^{(l)}(n) &= \sum_{j=1}^{\infty} \frac{1}{j!} \varepsilon(\varepsilon-1) \cdots (\varepsilon-j) \frac{1}{2} \{ [(l+1/2)^{\varepsilon-j} + [(l+1/2)^{\varepsilon-j}]^* \} n^j \\
 (49) \quad &= \sum_{j=1}^{\infty} \frac{1}{j!} \varepsilon(\varepsilon-1) \cdots (\varepsilon-j) |l+1/2|^{\varepsilon-j} \cos(\pi(\varepsilon-j))^{(1-sg(l))/2} n^j
 \end{aligned}$$

which is convergent for $|n/l| < 1$.

Next it is shown that,

Lemma 3.2.

$$(50) \quad \sum_n v_n^{(r)} w^{(l)}(n) = 0, \forall r, l$$

$\Downarrow$

$$(51) \quad \sum_n v_n^{(r)} n^m = 0 \quad m = 1, 2, \dots$$

Proof. This follows from the following argument. Multiplying (49) by $|l+1/2|^{1-\varepsilon}$ and taking the limit $l \rightarrow \infty$, leaves only the first term in (49) leading to,

$$(52) \quad \lim_{l \rightarrow \infty} |l+1/2|^{1-\varepsilon} w^{(l)}(n) = n$$

thus multiplying (50) by $|l+1/2|^{1-\varepsilon}$ and taking the limit implies,

$$(53) \quad \sum_n v_n^{(r)} n = 0$$

next multiplying (50) by $|l+1/2|^{2-\varepsilon}$, using (49), (53) and taking the limit $l \rightarrow \infty$ leads to,

$$(54) \quad \sum_n v_n^{(r)} n^2 = 0$$

continuing with this procedure leads to (51). $\square$

Replacement of the binomial expansion for $f_\alpha(n)$ in (35) and using (51) implies

$$(55) \quad \pi(d\omega) = 0$$

thus there are no non-trivial junk 1-forms for any of the two choices of $f_\alpha(k)$. $\square$

Similar, but lengthier, arguments lead to the absence of junk forms of any order.

The absence of junk forms has drastic consequences for the construction of the corresponding differential algebra. No quotient is needed. In the canonical case this quotient leads to the vanishing of any form of degree greater than one. In the deformed case there are forms of arbitrary degree and,

as can be seen from expression (32), the differential algebra is neither anti-commutative nor commutative. This constitutes a remarkable situation since starting from a commutative C^* -algebra in the triple, the deformed choice of the Dirac operator leads to a quite involved non-commutative differential algebra.

3.4. Distance. Let $\bar{\mathcal{A}}$ denote the closure of $\mathcal{A}$. A linear functional $\varphi : \bar{\mathcal{A}} \rightarrow \mathbb{C}$ is called a state. The set of all states being denoted by $S(\bar{\mathcal{A}})$. The distance $d(\varphi, \psi)$ between two states φ and ψ is defined by,

$$(56) \quad d(\varphi, \psi) = \sup |\varphi(a) - \psi(a)|$$

where the supremum is taken over all elements $a \in \bar{\mathcal{A}}$ such that,

$$(57) \quad \| [D, a] \| \leq 1$$

where $\| \cdot \|$ denotes the operator norm. The l.h.s. of (57) is given by,

$$(58) \quad \| [D, a] \| = \sup_{|\psi\rangle} \frac{|[D, a]|\psi\rangle|}{\| |\psi\rangle \|}$$

The vector whose norm appears in the r.h.s. of (58) is given by,

$$(59) \quad [D, a]|\psi\rangle = \sum_{l,m} a_{l-m} [f_{\alpha}(l) - f_{\alpha}(m)] \psi_m |l\rangle$$

where ψ_m denotes the Fourier component of $|\psi\rangle$ ($|\psi\rangle = \sum_m \psi_m |m\rangle$). The fact that for $\alpha \neq 0$ the quantity $f_{\alpha}(l) - f_{\alpha}(m)$ does not depend only on $l - m$ makes the evaluation of the norm of this vector for arbitrary $|\psi\rangle$ and $a(x)$ a non-trivial task. It is true however that,

$$(60) \quad f_{\alpha}(l) - f_{\alpha}(m) = f_{\alpha}(l - m) + O(\alpha)$$

therefore defining the following Liouville ε -derivative⁵(which reduces to the usual derivative for $\varepsilon = 1(\alpha = 0)$),

$$(61) \quad \frac{d^{\varepsilon}}{dx} g(x) = i \sum_l g_l f_{\alpha}(l) e^{-i(l+1/2)x}$$

(59) implies,

$$(62) \quad \begin{aligned} \langle x | [D, a] | \psi \rangle &= \sum_{l,m} a_{l-m} f(l - m) \psi_m \langle x | l \rangle + O(\alpha) \\ &= \psi(x) \left(-i \frac{d^{\varepsilon}}{dx} a(x) \right) + O(\alpha) \end{aligned}$$

thus,

$$(63) \quad \| [D, a] \| = \sup_{|\psi\rangle} \sqrt{\frac{\int dx |\psi(x)|^2 \left| \frac{d^{\varepsilon}}{dx} a(x) \right|^2}{\int dx |\psi(x)|^2}} + O(\alpha)$$

⁵This definition is an example of what is known in the literature as fractional derivative. See for example ref. [5] for an overview of the subject.

hence condition (57) implies,

$$(64) \quad \left| \frac{d^\epsilon}{dx} a(x) \right| \leq 1 + O(\alpha) \quad \forall x$$

Example 3.3. (i) *The canonical triple*[1] *over* S_1 .

In this case $f_\alpha(l) = l$ and the ε -derivative reduces to the usual derivative. Thus (57) leads to,

$$(65) \quad \left| \frac{d}{dx} a(x) \right| \leq 1 \quad \forall x$$

This last inequality implies that the functions $a(x)$ over which the supremum is taken are Lipschitz, i.e., they satisfy⁶,

$$(69) \quad |a(x_1) - a(x_0)| \leq |x_1 - x_0|$$

Next consider the state δ_{x_0} defined by,

$$(70) \quad \delta_{x_0}(a(x)) = a(x_0)$$

the distance between the two states δ_{x_1} and δ_{x_0} is, according to (56),

$$(71) \quad d(\delta_{x_1}, \delta_{x_0}) = \sup |\delta_{x_1}(a) - \delta_{x_0}(a)| = \sup |a(x_1) - a(x_0)|$$

thus (69) implies,

$$(72) \quad d(\delta_{x_1}, \delta_{x_0}) \leq |x_1 - x_0|$$

On the other hand, consider the distance function to a fixed point x_0 ,

$$(73) \quad d_{x_0}(x) = |x - x_0|$$

which is clearly a function satisfying,

$$(74) \quad \| [D, d_{x_0}] \| \leq 1$$

thus, it is clear that,

$$(75) \quad d(\delta_{x_1}, \delta_{x_0}) \geq |d_{x_0}(x_1) - d_{x_0}(x_0)| = |x_1 - x_0|$$

hence, (72) and (75) imply that,

$$(76) \quad d(\delta_{x_1}, \delta_{x_0}) = |x_1 - x_0|$$

(ii) $\alpha \neq 0$

⁶To see this note that (65) implies

$$(66) \quad \int_{x_0}^{x_1} dx \left| \frac{d}{dx} a(x) \right| \leq x_1 - x_0 \quad , x_1 > x_0$$

noting that,

$$(67) \quad \left| \int_{x_0}^{x_1} dx \frac{d}{dx} a(x) \right| \leq \int_{x_0}^{x_1} dx \left| \frac{d}{dx} a(x) \right|$$

eq. (69) is obtained, i.e.,

$$(68) \quad \left| \int_{x_0}^{x_1} dx \frac{d}{dx} a(x) \right| = |a(x_1) - a(x_0)| \leq |x_1 - x_0|$$

In what follows we neglect the $O(\alpha)$ corrections in (64). It turns out to be convenient to define the following kernel⁷,

$$\begin{aligned}
 d_{-\varepsilon}(x-y) &= \lim_{\eta \rightarrow 0} \sum_l \frac{1}{2 \cos(\pi\varepsilon)} \{[-(l+1/2-i\eta)]^{-\varepsilon} + c.c.\} e^{-i(l+1/2)(x-y)} \\
 &= \lim_{\eta \rightarrow 0} \sum_l |l+1/2-i\eta|^{-\varepsilon} [\cos(\pi\varepsilon)]^{\frac{sg(l)-1}{2}} e^{-i(l+1/2)(x-y)} \\
 (77) \quad &= \frac{1}{2 \cos(\pi\varepsilon)} \frac{|x|^{\varepsilon-1}}{\Gamma(\varepsilon)} e^{i\frac{\pi}{2}\varepsilon sg(x)}
 \end{aligned}$$

which has the following useful property,

$$\int_{-\pi}^{\pi} dy d_{-\varepsilon}(x-y) \frac{d^\varepsilon}{dy} a(y) = a(x)$$

and consider the following operator,

$$D_{-\varepsilon}^{\alpha\beta} g(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} dy |d_{-\varepsilon}(\beta-y) - d_{-\varepsilon}(\alpha-y)| g(y)$$

which is chosen to produce the following result,

$$\begin{aligned}
 D_{-\varepsilon}^{\alpha\beta} \left| \frac{d^\varepsilon}{dx} a(x) \right| &= \frac{1}{2\pi} \int_{-\pi}^{\pi} dy |d_{-\varepsilon}(\beta-y) - d_{-\varepsilon}(\alpha-y)| \frac{d^\varepsilon}{dy} a(y) \\
 &\geq \left| \frac{1}{2\pi} \int_{-\pi}^{\pi} dy (d_{-\varepsilon}(\beta-y) - d_{-\varepsilon}(\alpha-y)) \frac{d^\varepsilon}{dy} a(y) \right| \\
 &= |a(\beta) - a(\alpha)|
 \end{aligned}$$

therefore applying the operator $D_{-\varepsilon}^{\alpha\beta}$ to both sides of (64) implies,

$$|a(\beta) - a(\alpha)| \leq D_{-\varepsilon}^{\alpha\beta} 1$$

it is not difficult to see that (77) implies that $D_{-\varepsilon}^{\alpha\beta} 1$ depends only of $|\beta - \alpha|$, i.e.,

$$D_{-\varepsilon}^{\alpha\beta} 1 = \frac{D_{-\varepsilon}(|\beta - \alpha|)}{2\pi}$$

Using the last equality in (77) leads to the following expression for $D_{-\varepsilon}(x)$ in,

$$D_{-\varepsilon}(x) = \mathcal{F}(\varepsilon) \left\{ C(\varepsilon) |x|^\varepsilon + \frac{1}{\varepsilon} [(\pi+x)^\varepsilon + (\pi-x)^\varepsilon - 2\pi^\varepsilon] \right\}$$

⁷The last equality in eq.(77) is obtained replacing in the r.h.s. of the first equality the following integral representation of the fractional power of $l+1/2-i\eta$,

$$(l+1/2-i\eta)^{-\varepsilon} = \frac{i^\varepsilon}{\Gamma(\varepsilon)} \int_0^\infty d\tau \tau^{\varepsilon-1} e^{-i(l+1/2-i\eta)\tau}$$

Note the role played by the parameter η for the convergence of the integral in the r.h.s. of the last equation.

where,

$$\begin{aligned}\mathcal{F}(\varepsilon) &= |2\Gamma(\varepsilon) \cos(\pi\varepsilon)|^{-1} \\ C(\varepsilon) &= \int_0^1 dy \{ |1-y|^{\varepsilon-1} + |y|^{\varepsilon-1} - 2|1-y|^{\varepsilon-1} |y|^{\varepsilon-1} \cos(\pi\varepsilon) \}\end{aligned}$$

for the case $\varepsilon = 1$ these expressions give $\mathcal{F}(1) = 1/2$ and $C(1) = 2$ thus leading to the usual result, $D_{-1}(x) = |x|$.

To obtain the distance function one repeats the argument between eqs. (69) and (76) obtaining,

$$d(\delta_{x_1}, \delta_{x_0}) = \frac{D_{-\varepsilon}(|x_1 - x_0|)}{2\pi}$$




In Fig. 2,3 and 4 the function $D_{-\varepsilon}(x)$ is plotted for $\varepsilon = 1, 0.9$ and 0.8 respectively. According to subsection 3.2 these values of ε correspond to spaces with dimension $d = 1/\varepsilon = 1, 1.111\cdots, 1.25$. Including the $O(\alpha)$ corrections to (64) would produce corrections of $O(\alpha^2)$ to $D_{-\varepsilon}(x)$.

4. OUTLOOK

The results presented in this work may be considered as first steps in investigating these geometries. Many additional issues deserve to be studied. Some of them are: integration, operatorial structure of the differential algebra, metric volume element, gauge theories

5. ACKNOWLEDGMENTS

I want to express my gratitude to S. Capriotti, C. Fosco, S. Grillo and F. Mendez for valuable comments and suggestions.


 FIGURE 2. Distance for a space of dimension $d = 1$.

 FIGURE 3. Distance for a space of dimension $d = 1.11 \dots$.

 FIGURE 4. Distance for a space of dimension $d = 1.25$.

REFERENCES

- [1] A. Connes, *Noncommutative geometry*, Academic Press(1994).
- [2] See for example, J. Collins, *Renormalization*, Cambridge Univ. Press(1984).
- [3] A. Connes and H. Moscovici, *The local index formula in noncommutative geometry*, GAFA, **5**(1995)174.
- [4] F. L. Mendez, *Tripletes espectrales con dimensi3n no entera*, final work for the Licenciatura, adv. R.T., Instituto Balseiro(2000).

- [5] K. Oldham and J. Spannier, *The fractional calculus*, Academic Press(1974).

INSTITUTO BALSEIRO Y CENTRO ATÓMICO BARILOCHE.

E-mail address: `trinch@cab.cnea.gov.ar`